

Research Article

[Open Access](#) [Research Article](#) PTZAID:AUR-8-154

## Postnatal Outcomes in Cases of Prenatally Detected Dilated Fetal Bladder

Published On: October 30, 2024 | Pages: 028 - 034

Author(s): Jack Bocock, Panicos Shangaris\*, Massimo Garriboli and Srividhya Sankaran\*

Objectives: To explore the long-term outcomes in fetuses presenting with an enlarged bladder antenatally. Methods - Design: A retrospective cohort study was conducted over a 15-year period between 2003 and 2018. Setting and participants: Fetuses suspected to have enlarged bladders antenatally were included. Outcome measurements and statistical analysis: Analyses ...

[Abstract View](#) [Full Article View](#) DOI: 10.17352/aur.000054

[Open Access](#) [Research Article](#) PTZAID:AUR-8-153

## Association between Serum Cortisol Level and Erectile Function in Healthy Men from the General Population

Published On: October 15, 2024 | Pages: 023 - 027

Author(s): Babak Rezanezhad\*, Rasmus Borgquist and Saad Elzanaty

Purpose: The aim of the study is to investigate the association between serum levels of cortisol and erectile function in healthy men aged 45-60 years. Material and methods: The serum level of cortisol was measured. The international index of erectile function-5 items (IIEF-5) was used to assess the erectile function. Data were gathered on the age of participants, B ...

[Abstract View](#) [Full Article View](#) DOI: 10.17352/aur.000053

[Open Access](#) [Research Article](#) PTZAID:AUR-8-152

## Marital Status and Serum Level of Testosterone: In Relation to Serum Level of Cortisol

Published On: September 10, 2024 | Pages: 018 - 022

Author(s): Babak Rezanezhad\*, Rasmus Borgquist and Saad Elzanaty

Purpose: To explore the association between marital status and testosterone, and cortisol serum levels in 119 healthy men aged 45 to 60 years representing the general population. Material and Methods: Data on men's age, waist circumference, body mass index, smoking, alcohol consumption, and marital status were collected. Serum levels of LH, testosterone, and cortisol ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/aur.000052

## Case Report

[Open Access](#) | [Case Report](#) | PTZAID:AUR-8-151

### Bilateral Localised Primary Ureteric Amyloidosis

Published On: August 20, 2024 | Pages: 015 - 017

Author(s): Padmanabha Vijayan\*

Amyloidosis is a group of diseases in which insoluble proteins called amyloids deposit in tissues and cause problems with body functions. It is not a form of cancer but is a serious condition. It may be symptomatic or localized. The latter is a rare condition and the affection of the ureter has been reported in fewer than 50 cases so far. This case is unique because ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/aur.000051

[Open Access](#) | [Case Report](#) | PTZAID:AUR-8-150

### Pre-Sacral Neurofibroma Mistaken for a Uterine Fibroid

Published On: August 20, 2024 | Pages: 012 - 014

Author(s): Padmanabha Vijayan\*

A 33-year-old female patient who presented with urinary retention was found to have a large pelvic mass that was initially diagnosed as a pedunculated posterior wall uterine mass. Further evaluation revealed the mass to be a dumbbell-shaped neural tumor that extended into the sacral canal, corresponding to sacral segments 2, 3, and 4. The patient experienced no neurological ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/aur.000050