

In this issue

Research Article

[Open Access](#) [Research Article](#) PTZAID:AOR-6-231

Solving the problem of post-operative airway obstruction in Nasal/Sinus Surgery

Published On: September 15, 2020 | Pages: 091 - 097

Author(s): Robert Kotler*, Keith Wahl and Kimberly J Lee

Sinus surgery, septoplasty—with or without turbinate reduction—and rhinoplasty are among the most common surgical procedures performed by our specialty. In 2006, 600,000 sinus surgeries were performed in the United States [1]. A recent paper reported more than 300,000 rhinoplasties done per year [1,2]. Septoplasties and ancillary procedures accounted for an additional ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-1759.000131

[Open Access](#) [Research Article](#) PTZAID:AOR-6-229

Computed Tomography (CT) contributes in the functional outcome of Endoscopic Septoplasty

Published On: August 17, 2020 | Pages: 084 - 087

Author(s): Islam MA*, Mamoon TB, Miki FU, Chowdhury NH, Khan SR, Rahman ASL, Miah MAS and Mohammad T

Background and objective: Septoplasty is the common surgical procedure performed in nasal septal deviation. Nasal obstruction is the most frequent presentation to the otolaryngologist all over the world with septal deviation. Patients usually presented with headache, epistaxis, ophthalmological, otological, upper and lower respiratory tract infections; and many more. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-1759.000129

[Open Access](#) [Research Article](#) PTZAID:AOR-6-227

Non-Opioid Otolaryngology Medical Practice

Published On: July 28, 2020 | Pages: 076 - 079

Author(s): Samba Siva Reddy Bathula* and Dustin Jones DO

Introduction: Opioids have been used as pain medications in the United States since the late 1990s and have become increasingly common since then. Following the declaration of a public health emergency by the US Department of Health and Human Services (US-HHS), the author switched from opioid-based pain management to non-opioid-based pain management without any signif ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-1759.000127

Case Report

[Open Access](#) [Case Report](#) PTZAID:AOR-6-228

A Huge Para-pharyngeal Mass: Excision through Trans-cervical Approach

Published On: August 11, 2020 | Pages: 080 - 083

Author(s): Islam MA*, Mohammad T, Lutfur ASM, Chowdhury NH and Milki FU

Para-Pharyngeal Space (PPS) is an inverted cone-shaped potential space that extends from the hyoid bone to the base of the skull. It is filled with fat and areolar tissue lie laterally on either sides of pharynx; and is bounded and subdivided by various condensations of fascia. Tumors, although rare less than 1% of head neck neoplasm can arise from any structures of th ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-1759.000128

Use of ureteral catheter for under vision suctioning in rigid therapeutic bronchoscopy in children: A novel approach

Published On: July 27, 2020 | Pages: 071 - 075

Author(s): Krushnakumar V Kesan*, Sikha K Kesan, Arvind Ade and Paras Kothari

Background: Rigid bronchoscopy is a common procedure and treatment of choice for foreign body (FB) in respiratory tract. Blind suctioning using rigid suction cannula is associated with technical difficulties and complications. Under vision flexible cannula are expensive and not commonly available. This led us to discover the alternative of using Ureteral catheters fle ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-1759.000126

Laryngeal Adenoid Cystic Carcinoma: A rare neoplasm with still undefined treatment aspects. Case report and literature review

Published On: July 11, 2020 | Pages: 062 - 065

Author(s): Gianluigi Mariano Grilli, Rossana Giancaspro, Valeria Fiore, Anna Del Colle*, Luciano Enrico Renato Magaldi and Giuseppe Pannone

Adenoid Cystic Carcinoma (ACC) is a rare malignant tumor, accounting for about 1% of head and neck malignancies and 10% of salivary gland cancers. The laryngeal localization of ACC is particularly rare because of the paucity of accessory salivary glands in this anatomical district. Prevalence is highest in the fifth and sixth decade of life with a slight female predil ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-1759.000124

Case Series

The supraclavicular artery island flap in oral cavity reconstruction

Published On: August 25, 2020 | Pages: 088 - 090

Author(s): Khoi A Nguyen*, Nhat D Nguyen, Tam C Tran and Khuong H Le

Objective: To evaluate the advantages and disadvantages of the SAI flap as an alternative for free flap in reconstruction of oral cavity defects. Methods: We report a case series of 21 patients undergoing reconstructive operation from January 2018, through September 2019 after ablation of oral cavity malignant tumor. Results: Mean flap dimensions were 5.6 cm wide an ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-1759.000130

[Open Access](#) | [Case Series](#) | PTZAID:AOR-6-225

Natural facelift longevity: A unique observation in a deep plane facelift case series

Published On: July 14, 2020 | Pages: 066 - 070

Author(s): Brian P Dickinson* and Peter F Giacobazzi

Background: Facelift procedures are technically challenging and are difficult procedures to consistently achieve good results. Critical evaluation of one's facelift results in their practice over time is helpful to evaluate progress and to find areas to improve upon so that we can optimize patient results. We evaluated our charts and before and after photographs to im ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-1759.000125