

Research Article

[Open Access](#) [Research Article](#) PTZAID:ARDM-3-123

Effectiveness and Tolerability of Febuxostat vs Allopurinol in patients with Chronic Kidney Disease (CKD) on Conservative Therapy

Published On: June 19, 2017 | Pages: 030 - 033

Author(s): Filomena Panza, Chiara Ralli and Ennio Duranti*

Uric Acid appears to be one of the most important prognostic markers and one of the principal risk factors for cardiovascular diseases. The aim of our study was to assess three groups of patients: a first group treated with Allopurinol (AL), a second group not adequately controlled by treatment with AL or allergic to AL switched to treatment with Febuxostat (AL/FB) ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/2455-5495.000023](#)

[Open Access](#) [Research Article](#) PTZAID:ARDM-3-122

Factor causing late referral of CKD patients to Nephrology care

Published On: June 19, 2017 | Pages: 026 - 029

Author(s): Beena Salman, Muhammad Tahir, Ruqaya Qureshi, Murtaza Fakhruddin Dhrolia, Aasim Ahmad and Salman Imtiaz*

Introduction: Identification of the disease in its early period changes the outcome. Early reorganization of the chronic kidney disease and a timely referral to nephrologist also affects the prognosis of the disease. The factors which contribute in late referral are well known in the western population. There is a need to look into the factors in our population. ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/2455-5495.000022](#)

[Open Access](#) [Research Article](#) PTZAID:ARDM-3-121

Use of Interpreters for non-native English speaking Kidney Allograft Recipients and outcomes after Kidney Transplantation

Published On: June 01, 2017 | Pages: 020 - 025

Author(s): Sanna Tahir, Holly Gillott, Francesca Jackson Spence, Jay Nath, Jemma Mytton, Felicity Evison and Adnan Sharif*

Background: Language barriers are associated with worse health outcomes in the general population but data in kidney transplantation is lacking. This study tested the hypothesis that non-native English speakers using interpreters have poorer outcomes after kidney transplantation compared to native English speakers. Methods: A single-center retrospective study anal ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-5495.000021

[Open Access](#) [Research Article](#) PTZAID:ARDM-3-120

Quality of Life in Patients with Chronic Renal Failure and Some Affecting Factors

Published On: May 29, 2017 | Pages: 012 - 019

Author(s): Mukadder Mollaolu* and Gonca Deveci

Purpose: This study was conducted to determine the quality of life (QoL) in dialysis patients (DP) and the factors affecting the QoL. Material and Methods: 104 DP patients receiving treatment at a university hospital were included in this research, which was also a cross-sectional study. Data were collected with the Patient Information Form (PIF) and Kidney Diseas ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-5495.000020

[Open Access](#) [Research Article](#) PTZAID:ARDM-3-119

Encapsulating Peritoneal Sclerosis: Different clinical presentations and their management

Published On: May 19, 2017 | Pages: 007 - 011

Author(s): Jaime Manuel Justo-Janeiro (fi csfacs)*, Elizabeth Huerta-Calixto, de la Rosa Paredes Rene, Fernando Vazquez de Lara and Luis G Vazquez de Lara

Introduction: One of most common treatments of end-stage renal disease is peritoneal dialysis. Encapsulating peritoneal sclerosis is a complication in which the osmotic capacity of the peritoneal barrier is lost, due to infections or the irritating effect of dialysis solutions. This pathology has different clinical presentations, hence the need of different diagnos ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-5495.000019

Short-Term Effects of Pre/Probiotics on P-Cresol and Indoxyl-Sulphate Serum Concentrations During the Various Stages of Chronic Kidney Disease

Published On: May 03, 2017 | Pages: 001 - 005

Author(s): Filomena Panza, Diletta Duranti, Ralli Chiara, Matteo Basile, Marco Bagnati, Giorgio Bellomo and Ennio Duranti*

The uremic syndrome is provoked by a progressive number of compounds that are normally excreted by kidneys in healthy individuals. Indoxylsulphate (IXS) and p-cresylsulphate (PCS), have been found increased in subjects with end stage renal disease (ESRD) creating great harm to biological systems; these uremic toxins come from the intestinal bacterial fermentation of t ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-5495.000017

Letter to Editor

Nephrolithiasis in hypercalcemic and normocalcemic primary Hyperparathyroidism

Published On: May 17, 2017 | Pages: 006 - 006

Author(s): Vitorino Modesto dos Santos*

Primary hyperparathyroidism (PHPT) is a common condition affecting up to 4 per 1000 of population, and the majority of cases are due to adenoma or hyperplasia of the gland [1]. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-5495.000018